

University of Groningen

**Erratum: elemental abundances in Milky Way-like galaxies from a hierarchical galaxy formation model**

De Lucia, Gabriella; Tornatore, Luca; Frenk, Carlos S.; Helmi, Amina; Navarro, Julio F.; White, Simon D. M.

*Published in:*  
Monthly Notices of the Royal Astronomical Society

*DOI:*  
[10.1093/mnras/stu2609](https://doi.org/10.1093/mnras/stu2609)

**IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.**

*Document Version*  
Publisher's PDF, also known as Version of record

*Publication date:*  
2015

[Link to publication in University of Groningen/UMCG research database](#)

*Citation for published version (APA):*

De Lucia, G., Tornatore, L., Frenk, C. S., Helmi, A., Navarro, J. F., & White, S. D. M. (2015). Erratum: elemental abundances in Milky Way-like galaxies from a hierarchical galaxy formation model. *Monthly Notices of the Royal Astronomical Society*, 447(4), 3420-3421. <https://doi.org/10.1093/mnras/stu2609>

**Copyright**

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

**Take-down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

# Erratum: elemental abundances in Milky Way-like galaxies from a hierarchical galaxy formation model

by Gabriella De Lucia,<sup>1</sup>★ Luca Tornatore,<sup>1</sup> Carlos S. Frenk,<sup>2</sup> Amina Helmi,<sup>3</sup> Julio F. Navarro<sup>4</sup> and Simon D. M. White<sup>5</sup>

<sup>1</sup>*INAF, Astronomical Observatory of Trieste, via G.B. Tiepolo 11, I-34143 Trieste, Italy*

<sup>2</sup>*Institute of Computational Cosmology, University of Durham, Science Laboratories, South Road, Durham DH13LE, UK*

<sup>3</sup>*Kapteyn Astronomical Institute, University of Groningen, PO Box 800, NL-9700 AV Groningen, the Netherlands*

<sup>4</sup>*Department of Physics and Astronomy, University of Victoria, Victoria, BC V8P 5C2, Canada*

<sup>5</sup>*Max Planck Institut für Astrophysik Karl-Schwarzschild-Str. 1, D-85741 Garching, Germany*

**Key words:** errata, addenda – Galaxy: abundances – Galaxy: evolution – Galaxy: formation – galaxies: dwarf.


Our paper ‘Elemental abundances in Milky Way-like galaxies from a hierarchical galaxy formation model’ was published in MNRAS, 445, 970 (2014). Due to a mistake, the code used for the runs included in the paper did not account correctly for the decrease of the galaxy stellar mass, following restitution of gas to the interstellar medium through stellar winds and supernovae explosions. As a consequence, the stellar masses given in the paper were slightly

overpredicted. This affects in particular Figs 7 and 9 of our original manuscript. The updated versions of these figures are reproduced here for completeness (see Figs 1 and 2).


For Aq-B-2, the predicted stellar mass of the model Milky Way galaxy is  $4.5 \times 10^{10} M_{\odot}$  (against  $\sim 6 \times 10^{10} M_{\odot}$  quoted in our paper), that is still in very good agreement with the estimated value for our own Milky Way. The current level of star formation rate is  $\sim 3.6 M_{\odot} \text{ yr}^{-1}$  ( $\sim 2.8 M_{\odot} \text{ yr}^{-1}$  in our paper), and the present value of the SNIa rate is  $\sim 0.73$  (0.58 in our paper) events/century, still higher than a factor of 2 to 3 with respect to the observational estimate of  $\sim 0.2$ – $0.3$  events/century.

All conclusions of our paper are unaffected.

\* E-mail: [delucia@oats.inaf.it](mailto:delucia@oats.inaf.it)



**Figure 1.** Physical properties of our model Milky Way galaxies, for the different simulations used in this study. The open circles show results from the model based on the instantaneous recycling approximation, while filled symbols show the corresponding results based on the updated chemical model presented in our paper. The red horizontal lines in each panel indicate observational estimates.



**Figure 2.** Star formation history for the model Milky Way galaxy in the run Aq-A-3. The solid black line shows the prediction from the reference model used in our study, while the black dashed line shows the corresponding prediction from a run that adopts an instantaneous recycling approximation. The coloured dotted, dot-dashed and long-dashed lines correspond to variations of the latter run with varying parameters.

This paper has been typeset from a  $\text{\LaTeX}$  file prepared by the author.